

2015 Algebra Prelim

September 14, 2015

INSTRUCTIONS: Do as many of the eight problems as you can. Four completely correct solutions will be a pass; a few complete solutions will count more than many partial solutions. Always carefully justify your answers. If you skip a step or omit some details in a proof, point out the gap and, if possible, indicate what would be required to fill it in.

- Find an irreducible polynomial of degree 5 over the field \mathbb{Z}_2 of two elements and use it to construct a field of order 32 as a quotient of the polynomial ring $\mathbb{Z}_2[x]$.
 - Using the polynomial you found in part (a), find a 5×5 matrix M over \mathbb{Z}_2 of order 31, so that $M^{31} = I$ but $M \neq I$.
- Find the minimal polynomial of $\sqrt{2} + \sqrt{3}$ over \mathbb{Q} . Justify your answer.
 - Let R be a commutative ring with no nonzero nilpotent elements. Show that the only units in the polynomial ring $R[x]$ are the units of R , regarded as constant polynomials.
 - Find all units in the polynomial ring $\mathbb{Z}_4[x]$.
- Let p and q be two distinct primes. Prove that there is at most one non-abelian group of order pq (up to isomorphisms) and describe the pairs (p, q) such that there is no non-abelian group of order pq .
 - Let L be a Galois extension of a field K of degree 4. What is the minimum number of subfields there could be strictly between K and L ? What is the maximum number of such subfields? Give examples where these bounds are attained.
 - How do these numbers change if we assume only that L is separable (but not necessarily Galois) over K ?
- Let R be a commutative algebra over \mathbb{C} . A derivation of R is a \mathbb{C} -linear map $D : R \rightarrow R$ such that (i) $D(1) = 0$, and (ii) $D(ab) = D(a)b + aD(b)$ for all $a, b \in R$.
 - Describe all derivations of the polynomial ring $\mathbb{C}[x]$.
 - Let A be the subring (or \mathbb{C} -subalgebra) of $\text{End}_{\mathbb{C}}(\mathbb{C}[x])$ generated by all derivations of $\mathbb{C}[x]$ and the left multiplications by x . Prove that $\mathbb{C}[x]$ is a simple left A -module. Note that the inclusion $A \rightarrow \text{End}_{\mathbb{C}}(\mathbb{C}[x])$ defines a natural left A -module structure on $\mathbb{C}[x]$.
 - Let G be a non-abelian group of order p^3 with p a prime.
 - Determine the order of the center Z of G .
 - Determine the number of inequivalent complex 1-dimensional representations of G .
 - Compute the dimensions of all the inequivalent irreducible representations of G and verify that the number of such representations equals the number of conjugacy classes of G .
- Prove that every finitely generated projective module over a commutative noetherian local ring is free.